REACTION
0000000 APPEARANCE
DDDD CAUSATIVED DRUGS
Exanthema
Maculopapular(flat/slightly raised in pimples), scarlatiniform/rubelliform(red) pruritic(itchy) lesion
Benzodiazepines (BZDs) Carbamazepine, Valproate, Lamotrigine
Urticaria
Erythematous(red), pruritic(itchy) wheals. May progress to angio-oedema and anaphylaxis
Benzodiazepines (BZDs) Carbamazepine, Valproate
Fixed drug eruptions
Reddened round lesions which blister and crust. Hyperpigmentation may be permanent.
Lesions tend to present in same location if rechallenge
performed
BZDs and carbamazepine
Bullous reactions
Blisters and vesicles; may be part of more generalised CDR such as vasculitis or EM complex*
Carbamazepine
Lichenoid reactions
Flat violaceous papules resembling lichen planus
Carbamazepine
Alopecia
Hair loss,Thinning hair
Carbamazepine, vigabatrin, valproate, BZDs
Pigmentation changes
Colour changes may affect skin, hair, nails, mucous membranes; may be permanent
BZDs, Valproate
Photosensitivity
Abnormal sensitivity to UVA or UVB (photoallergy); may result in severe sunburn
Carbamazepine ; BZDs
Exfoliative dermatitis
Generalised redness, pruritus and eczema with exfoliation(skin peeling off); may affect whole bo
Carbamazepine, diazepam , valproate

Carbamazepine

Vasculitis
Small haemorrhagic macules coalescing to larger blisters and ulcers; fever, malaise, joint/musc
BZDs, Carbamazepine, Valproate
Anticonvulsant hypersensitivity syndrome
Multiorgan syndrome affecting skin (exanthematous pustulation), fever, lymphadenopathy, leuc
Erythema multiforme (EM) complex*
Sudden onset skin eruption(morbilliform) with□ stomatitis, vesicles around lips and oral mucos
Carbamazepine, valproate
Erythema nodosum
Painful erythematous subcutaneous nodules: fever malaise: general symptoms may persist for

SJS= Stevens- Johnson Syndrome; TEN= Toxic Epidermal Necrolysis