REACTION

Exanthema

Maculopapular(flat/slightly raised in pimples), scarlatiniform/rubelliform(red) pruritic(itchy) lesion Benzodiazepines (BZDs) Carbamazepine, Valproate, Lamotrigine

Urticaria

Erythematous(red), pruritic(itchy) wheals. May progress to angio-oedema and anaphylaxis Benzodiazepines (BZDs) Carbamazepine, Valproate

Fixed drug eruptions Reddened round lesions which blister and crust. Hyperpigmentation may be permanent.

Lesions tend to present in same location if rechallenge performed BZDs and carbamazepine

Bullous reactions Blisters and vesicles; may be part of more generalised CDR such as vasculitis or EM complex* Carbamazepine

Lichenoid reactions Flat violaceous papules resembling lichen planus Carbamazepine

Alopecia Hair loss,Thinning hair Carbamazepine, vigabatrin, valproate, BZDs

Pigmentation changes Colour changes may affect skin, hair, nails, mucous membranes; may be permanent BZDs, Valproate

Photosensitivity Abnormal sensitivity to UVA or Carbamazepine ; BZDs

Exfoliative dermatitis Generalised redness, pruritus and eczema with exfoliation(skin peeling off); may affect whole bo Carbamazepine, diazepam , valproate

Vasculitis

Small haemorrhagic macules coalescing to larger blisters and ulcers; fever, malaise, joint/muscl BZDs, Carbamazepine, Valproate

Anticonvulsant hypersensitivity syndrome Multiorgan syndrome affecting skin (exanthematous pustulation), fever, lymphadenopathy, leuco

Erythema multiforme (EM) complex*

Sudden onset skin eruption(morbilliform) with stomatitis, vesicles around lips and oral mucosa Carbamazepine, valproate

Erythema nodosum

Painful erythematous subcutaneous nodules; fever, malaise; general symptoms may persist for Carbamazepine

SJS= Stevens- Johnson Syndrome; TEN= Toxic Epidermal Necrolysis